

only if each second repeat test animal meets the requirements prescribed in paragraph (d) of this section.

(f) [Reserved]

(g) *Exceptions*—(1) The test prescribed in this section need not be performed for Whole Blood, Red Blood Cells, Cryoprecipitated AHF, Platelets, Plasma, or Cellular Therapy Products.

(2) [Reserved]

[41 FR 10891, Mar. 15, 1976, as amended at 49 FR 15187, Apr. 18, 1984; 49 FR 23834, June 8, 1984; 50 FR 4133, Jan. 29, 1985; 51 FR 15607, Apr. 25, 1986; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994; 63 FR 19403, Apr. 20, 1998; 63 FR 41718, Aug. 5, 1998]

#### § 610.11a Inactivated influenza vaccine, general safety test.

For inactivated influenza vaccine, the general safety test shall be conducted in the manner indicated in § 610.11 of this chapter except that, with reference to guinea pigs, the test shall be satisfied if the product provides satisfactory results using either the subcutaneous or intraperitoneal injection of 5.0 milliliters of inactivated influenza vaccine into each guinea pig. The requirements for general safety for inactivated influenza vaccine shall not be considered to be satisfied unless each lot of influenza vaccine is assayed for endotoxin in comparison to a reference preparation provided by the Food and Drug Administration, and such lot is found to contain no more endotoxin than the reference preparation.

[39 FR 40016, Nov. 13, 1974]

#### § 610.12 Sterility.

Except as provided in paragraphs (f) and (g) of this section, the sterility of each lot of each product shall be demonstrated by the performance of the tests prescribed in paragraphs (a) and (b) of this section for both bulk and final container material.

(a) *The test.* Bulk material shall be tested separately from final container material and material from each final container shall be tested in individual test vessels as follows:

(1) *Using Fluid Thioglycollate Medium*—(i) *Bulk and final container material.* The volume of product, as required by paragraph (d) of this section (hereinafter referred to also as the

“inoculum”), from samples of both bulk and final container material, shall be inoculated into test vessels of Fluid Thioglycollate Medium. The inoculum and medium shall be mixed thoroughly and incubated at a temperature of 30 to 35 °C for a test period of no less than 14 days and examined visually for evidence of growth on the third, fourth, or fifth day, and on the seventh or eighth day, and on the last day of the test period. Results of each examination shall be recorded. If the inoculum renders the medium turbid so that the absence of growth cannot be determined reliably by visual examination, portions of this turbid medium in amounts of no less than 1.0 milliliter shall be transferred on the third, fourth, or fifth day of incubation, from each of the test vessels and inoculated into additional vessels of the medium. The material in the additional vessels shall be incubated at a temperature of 30 to 35 °C for no less than 14 days. Notwithstanding such transfer of material, examination of the original vessels shall be continued as prescribed above. The additional test vessels shall be examined visually for evidence of growth on the third, fourth, or fifth day of incubation, and on the seventh or eighth day, and on the last day of the incubation period. If growth appears, repeat tests may be performed as prescribed in paragraph (b) of this section and interpreted as specified in paragraph (c) of this section.

(ii) *Final container material containing a mercurial preservative.* In addition to the test prescribed in paragraph (a)(1)(i) of this section, final container material containing a mercurial preservative shall be tested using Fluid Thioglycollate Medium following the procedures prescribed in such subparagraph, except that the incubation shall be at a temperature of 20 to 25 °C.

(2) *Using Soybean-Casein Digest Medium.* Except for products containing a mercurial preservative, a test shall be made on final container material, following the procedures prescribed in paragraph (a)(1)(i) of this section, except that the medium shall be Soybean-Casein Digest Medium and the incubation shall be at a temperature of 20 to 25 °C.